

Tennessee Behavioral Risk Factor Surveillance Survey 2012 State and Regional Weighted Data

Have you ever been told by a doctor, nurse, or other health professional that you had asthma?

| State and Regions | Total Respondents | <i>Diagnosed With Asthma</i> ¹ | <i>Weighted Risk</i> ² | <i>Population at Risk</i> ³ |
|-------------------------------|-------------------|---|-----------------------------------|--|
| TENNESSEE ⁴ | 7,036 | 736 | 11.0 | 545,592 |
| Northeast | 443 | 56 | 10.7 | 29,298 |
| East | 634 | 76 | 12.3 | 72,431 |
| Southeast | 505 | 56 | 13.5 | 34,172 |
| Upper Cumberland | 459 | 61 | 12.2 | 32,378 |
| Mid-Cumberland | 753 | 80 | 10.6 | 89,280 |
| South Central | 494 | 41 | 6.6 | 19,269 |
| Northwest | 392 | 38 | 13.5 | 26,513 |
| Southwest | 442 | 49 | 13.8 | 29,856 |
| Shelby | 519 | 50 | 10.4 | 72,326 |
| Davidson | 559 | 54 | 7.8 | 39,333 |
| Knox | 518 | 60 | 11.9 | 40,897 |
| Hamilton | 458 | 44 | 7.9 | 21,269 |
| Madison | 443 | 25 | 4.2 | 3,105 |
| Sullivan | 387 | 44 | 8.9 | 10,772 |

¹ Number of respondents who answered YES.

² Weighted risk is the estimated percentage of adult population who have ever had asthma based on the survey.

³ Population at risk is the estimated population who have ever had asthma based on the survey. Note: The sum of regional estimates in this column will not add up to statewide total.

⁴ Statewide estimates include additional survey coverage by cell phone and different weighting methodology.