



# Let's Talk About Hemorrhagic Strokes and Their Causes

## Did I have a hemorrhagic stroke?


Not all strokes are caused by blood clots that block an artery. Fifteen to 20 percent happen when a blood vessel ruptures in or near the brain. This is called a *hemorrhagic* (hem-o-RAYG-ik) stroke. (See figure.)

You may have many diagnostic tests before your doctor determines that you had a hemorrhagic stroke. Hemorrhagic strokes are different from ischemic strokes in many ways.

- The fatality rate is higher and overall prognosis poorer for those who have hemorrhagic strokes.
- People who have hemorrhagic strokes are younger.
- This kind of stroke is often associated with a very severe headache, nausea and vomiting.

- Usually the symptoms appear suddenly.
- A transient ischemic attack (TIA) or any other stroke warning sign may not precede this type of stroke.

Ruptured artery in the brain stops oxygen supply to the surrounding areas of the brain.



A hemorrhagic stroke can occur when an aneurysm (a blood-filled pouch that balloons out from an artery) ruptures, flooding the surrounding tissue with blood.

## Are all hemorrhagic strokes the same?

There are two kinds of hemorrhagic stroke. In both, a blood vessel ruptures, disrupting blood flow to part of the brain. Increasing the risk of hemorrhagic strokes are: 1) cigarette smoking; 2) use of oral contraceptives (particularly those with high estrogen content); 3) excessive alcohol intake; and 4) use of illegal drugs.

### **Subarachnoid** (sub-ah-RAK-noid) **hemorrhage:**

- Occurs when a blood vessel on the surface of the brain ruptures and bleeds into the space between the brain and the skull.

- A ruptured aneurysm, often caused by high blood pressure, is the most common cause. An aneurysm is a blood-filled pouch that balloons out from an artery wall.

### **Intracerebral hemorrhage:**

- Occurs when a blood vessel bleeds into the tissue deep within the brain. This type of hemorrhage is responsible for about 10 percent of all strokes.
- Chronically high blood pressure or aging blood vessels are the main causes of this type of stroke.

## How are hemorrhagic strokes treated?

Because hemorrhages may be life-threatening, hospital care is required in an Intensive Care Unit. Medication can control further bleeding as well as control high blood pressure. Other medicine may be prescribed

to reduce the brain swelling that follows a stroke. Surgery may be needed depending on the cause of the hemorrhage. Surgery could be done to repair an aneurysm or remove a blood clot.

## How can I learn more?

- Talk to your doctor, nurse or other healthcare professionals. Ask about other stroke topics. This is one of many *Let's Talk About Stroke* fact sheets available.
- For more information on stroke, or to receive additional fact sheets, call the American Stroke Association at 1-888-4-STROKE (1-888-478-7653) or visit us online at [StrokeAssociation.org](http://StrokeAssociation.org).
- If you or someone you know has had a stroke, call the American

Stroke Association's "Warmline" at 1-888-4-STROKE (1-888-478-7653), and:

- ✓ Speak with other stroke survivors and caregivers, trained to answer your questions and offer support
- ✓ Get information on stroke support groups in your area
- ✓ Sign up to get *Stroke Connection Magazine*, a free publication for stroke survivors and caregivers

## What are the Warning Signs of Stroke?

-  **Sudden weakness or numbness of the face, arm or leg, especially on one side of the body**
-  **Sudden confusion, trouble speaking or understanding**
-  **Sudden trouble seeing in one or both eyes**
-  **Sudden trouble walking, dizziness, loss of balance or coordination**
-  **Sudden, severe headaches with no known cause**



Learn to recognize a stroke.  
*Because time lost is brain lost.*

Today there are treatments that can reduce the risk of damage from the most common type of stroke, but only if you get help quickly – within 3 hours of your first symptoms.

Call 9-1-1 immediately if you experience these warning signs!

## Do you have questions for your doctor or nurse?

Take a few minutes to write your own questions for the next time you see your healthcare provider:

Will I have to worry about a hemorrhagic stroke occurring again?  
\_\_\_\_\_

What can I do to help prevent another stroke?  
\_\_\_\_\_

How can I control high blood pressure?  
\_\_\_\_\_

What else should I know about treatment options?  
\_\_\_\_\_  
\_\_\_\_\_

*The statistics in this kit were up to date at publication. For the latest statistics, see the Heart Disease and Stroke Statistics Update at [americanheart.org/statistics](http://americanheart.org/statistics).*

*The American Stroke Association is a division of the American Heart Association. Your contributions will support research and educational programs that help reduce disability and death from stroke.*

©1999-2003, American Heart Association.

**American Stroke  
Association**<sup>SM</sup>

A Division of American  
Heart Association

