


## Neonatal Abstinence Syndrome (NAS) among TennCare Enrollees: 2021 data

Neonatal Abstinence Syndrome is a withdrawal syndrome experienced by newborn infants shortly after birth when they were exposed to certain substances, most often opioids, causing dependence during pregnancy. The infant is exposed in the womb before birth when a mother uses substances such as prescriptions medications or illicit drugs during pregnancy. After birth, the newborn infant can go through withdrawal once the infant is no longer exposed to or receives these substances. Infants with NAS may stay in the hospital longer than other babies to treat these symptoms.

This report documents the occurrence of neonatal abstinence syndrome among TennCare enrollees in 2021. Cases of neonatal abstinence syndrome (NAS) were identified based on the presence of ICD10 diagnosis codes P96.1<sup>1</sup> and P96.2<sup>1</sup> transmitted to TennCare from medical providers billing for services provided to infants during the first year of life. This study included infants born between January 1 and December 31, 2021; where other years are provided for comparison purposes, those cohorts were born during the specified calendar year. TennCare eligibility status was determined using TennCare’s Interchange system. Cases were identified from infants that were eligible at time of birth or enrolled in TennCare during their first year of life. The number of live births, used as the denominator, were determined based on a linkage of vital statistics records and TennCare eligibility records.

**Figure 1: Incidence of Neonatal Abstinence Syndrome among TennCare Enrollees**



As Figure 1 illustrates, there was an increase in the incidence rates of NAS per 1,000 live births among TennCare recipients from CY2012 to CY2016. However, the rate of NAS births has decreased for several successive years and remained stable over the past three years.


<sup>1</sup> Definition: Drug withdrawal syndrome in a newborn, excluding fetal alcohol syndrome.

**Table 1: NAS Mother’s TennCare Status at Time of Delivery**

Calendar Year	TennCare Newborns Treated for NAS During Year	Mothers on TennCare at Time of TennCare NAS Birth	Percent of TennCare NAS Infants Born to TennCare Mothers	Mothers NOT on TennCare at Time of TennCare NAS Birth	Percent of TennCare NAS Infants NOT Born to TennCare Mothers
2008	264	229	87%	35	13%
2009	444	335	75%	109	25%
2010	512	424	83%	88	17%
2011	528	483	91%	45	9%
2012	736	613	83%	123	17%
2013	943	823	87%	120	13%
2014	1,101	1,017	92%	84	8%
2015	1,197	1,098	92%	99	8%
2016	1,357	1,261	93%	96	7%
2017	1,363	1,254	92%	109	8%
2018	1,181	1,093	93%	88	7%
2019	992	922	93%	70	7%
2020	957	890	93%	67	7%
2021	989	919	93%	70	7%


Table 1 presents information regarding the TennCare status of mothers of TennCare NAS infants at the time of birth. In 2021, 93% of TennCare NAS infants were born to mothers who were eligible in TennCare at the time of delivery, while the remaining 7% of NAS infants were born to mothers who were not TennCare-eligible at the time of delivery. This ratio has remained steady since 2014.

**Figure 2: Eligibility Span for TennCare NAS Mothers through Pregnancy Period**







**Figure 3: Demographic Characteristics of NAS Mothers**



**Figure 3A**



**Figure 3B**



**Figure 3C**

Figures 3A, 3B, and 3C display basic demographic characteristics of NAS mothers who had TennCare eligibility coverage in CY2021. White Non-Hispanic mothers have the highest NAS rate (3.4%) among all the races (see Figure 3A). As displayed in Figure 3B, while women in the 25–29-year-old age category have delivered more NAS babies than any other age group, women in the 35–39-year-old age category have the highest rate of NAS deliveries. Generally, the risk of a baby being delivered with neonatal abstinence syndrome increases with the age of the mother. Figure 3C shows the delivery rate of NAS mothers in East Tennessee is 3.1% which is higher than Middle (2.4%) and West Tennessee (1.1%).

**Table 2: Percentage of Infants in DCS Custody within One Year of Birth**

Metric	All Infants	NAS Infants
Total Number of Infants	49,333	992
Number of Infants in DCS Custody	549	233
Percent of Infants in DCS Custody	1.1%	23.5%

Using TennCare eligibility records, it was determined that 233 of the 992 infants diagnosed with NAS in CY2021 (23.5%) were placed in Department of Children’s Services (DCS) custody within one year of their birth. Among all TennCare infants born in CY2021, 1.1% were placed in DCS within one year of birth (Table


2). Infants born with NAS are about 21.1 times more likely to be in DCS custody during their first year of life as compared to other TennCare infants.

**Table 3: Impact of NAS on Infant Health Care Expenditures<sup>2</sup>**

Metric	All TennCare paid live births	All TennCare normal birth weight births	All TennCare live low birth weight births	NAS Infants
Number of Births	45,713	40,605	5,108	1,001
Total Cost for Infants in First Year of Life	\$518,062,955	\$272,431,834	\$245,631,121	\$52,525,797
Average Cost per Child	\$11,333	\$6,709	\$48,088	\$52,473
Average length of Stay	5.2	3.3	20.5	29.2


To determine the financial impact of NAS relative to all births, TennCare’s Interchange System was used to quantify expenditures for live born infants in the first year of life (Table 3). In CY2021, the average cost of care for a NAS infant in the first year of life is more than 7.8 times higher than the average cost of care for normal birth weight infants and more than 1.1 times higher than the average cost of care for low-birth-weight infants.

**Figure 4: Number and Percentage of Unique Opioid Types for NAS Mothers**



<sup>2</sup> Includes all expenditures paid through the first year of life. Totals are subject to change based on updated data.

**Figure 5: Number and Percentage of Unique Prescribers for NAS Mothers**



All opioid claims up to one year prior to birth for any woman with a NAS infant were evaluated. Medication Assisted Treatment medications, which are an evidence-based treatment and best-practice of care for members with opioid use disorder, are also considered opioids and were included in this analysis. Figures 4 and 5<sup>3</sup> illustrate the numbers of NAS mothers were using different TennCare-paid opioid prescriptions as well as the number of unique opioid types during the pregnancy period. Overall, a total of 4,925 TennCare-paid prescriptions for opioids were issued to 440 women with NAS infants. The overall percentage of mothers who did not have any opioid prescriptions paid for by TennCare was 50.6% (450/890) in CY2020. This value was similar to the number observed in CY2019 of 49.1% (469/922). Among NAS mothers with at least one opioid prescription paid by TennCare, each mother had on average 11.19 opioid prescriptions and from the remaining 49.4% women who received one opioid prescription, 66% of NAS mothers were received one or two types of opioids in the year of pregnancy period.

All opioid claims up to one year prior to birth for any woman with a NAS infant were evaluated. Medication Assisted Treatment (MAT) medications, which is an evidence-based treatment and best-practice of care for members with opioid use disorder, are also considered opioids and were included in this analysis.

<sup>3</sup> Any pharmacy claim with a National Drug Code correlation to the following Therapeutic Class Codes (HIC3) was considered an opioid: H3A, H3H, H3J, H3M, H3N, H3R, H3T, H3U, H3W or H3X.

Figures 4 and 5<sup>4</sup> illustrate the numbers of NAS mothers were using different TennCare-paid opioid prescriptions as well as the number of unique opioid types during the pregnancy period. The overall percentage of mothers who did not have any opioid prescriptions paid for by TennCare was 50.6% (465/919) in CY2021. This value was similar to the number observed in CY2020 of 50.6% (450/890). Of the women who received an opioid prescription paid for by TennCare, over 45% were receiving medication assisted treatment for opioid or substance use disorder. Medication-assisted treatment (MAT) is the use of FDA approved medications for treatment of opioid use disorder in combination with counseling and behavioral therapies. MAT works by reducing cravings and controlling withdrawal symptoms. MAT can also reduce or eliminate polysubstance use, decreases illicit opioid use, and improve maternal and infant outcomes. Individuals who receive medication assisted treatment and recovery services are more likely to have a full-term birth and less likely to have a low-birth weight baby. Note, this data does not account for services provided in an institutional setting, such as an inpatient hospital, or other forms of addiction treatment where a separate pharmacy claim does not exist. Overall, these results show continued improvement in reducing opioid utilization and improving access to Medication Assisted Treatment and recover services for opioid use disorder. Both interventions have been integral to reducing the overall number and severity of NAS births for TennCare members.

**Table 4: Narcotic Analgesic and Contraceptive Use Among All TennCare Women**

Demographics (Years)	TennCare Women	Women Prescribed Opioid (>30 days supplied)	Opioid Users Rate per 1,000	Women Prescribed Contraceptives and Opioid	% of Women on Opioid and Contraceptives	Women Prescribed Opioid without Contraceptives	% of Women on Opioid without Contraceptives
15 - 20	113,101	114	1	41	36%	73	64%
21 - 24	55,741	460	8	119	26%	341	74%
25 - 29	72,937	1,846	25	379	21%	1,467	79%
30 - 34	75,033	3,028	40	493	16%	2,535	84%
35 - 39	62,526	3,619	58	390	11%	3,229	89%
40 - 44	48,629	3,506	72	259	7%	3,247	93%
<b>Total</b>	<b>427,967</b>	<b>12,573</b>	<b>29</b>	<b>1,681</b>	<b>13%</b>	<b>10,892</b>	<b>87%</b>

Note: This analysis underestimates use of voluntary long-acting reversible contraceptives (vLARC). Additionally, this metric does not account for individuals with permanent forms of contraception such as tubal ligation or hysterectomy.

The rate of women using prescribed opioids and contraceptive medications was determined in CY2021 (Table 4). The analysis was limited only to women of child-bearing age (15–44). The prescription histories of TennCare women of child-bearing age were evaluated for the presence of opioids and contraceptive products<sup>5</sup>. Pharmacy claims were used in this analysis and thus the claims more readily capture prescribed contraceptive options (e.g. oral, intrauterine devices, injectable, transdermal, hormonal ring). Due to


<sup>4</sup> Any pharmacy claim with a National Drug Code correlation to the following Therapeutic Class Codes (HIC3) was considered an opioid: H3A, H3H, H3J, H3M, H3N, H3R, H3T, H3U, H3W or H3X.

<sup>5</sup> Any pharmacy claims with an NDC correlating to any HIC3 codes of G8A, G8B, G8C, G8F, G9B, X1C, G9A was considered a contraceptive.



limitations in pharmacy claims, this analysis does not capture all women who may have used a long-acting reversible contraceptive (LARC) and those who have permanent sterilization. Thus, this table is an underrepresentation of the full contraceptive options available to and being utilized by TennCare mothers. Women were excluded from the analysis if they had opioid prescriptions totaling less than 30 days in CY2021. As Table 4 indicates, approximately 13% of women of child-bearing age in 2021 who were prescribed opioids for more than 30 days in a year also received a form of contraceptive.

**Figure 6: Opioid and Contraceptives Trends for TennCare Women**



Based on the 5-year utilization data of opioid and contraceptives among TennCare women aged 15 to 44 years old, the rate of opioid users per 1,000 women continuously decreased from 2016-2019 and slightly increased in 2020 and decreased in 2021. In 2021, the overall rate of prescription opioid utilization among women aged 15-44 was 29.4 opioid users per 1,000 eligible women, a 11% decrease compared to 2020 (see Figure 6 below). Similar to Table 4, this does not capture all individuals with contraception, particularly LARC and permanent sterilization. Figure 6 shows 74% decrease in the rate of opioid use among women aged 15-44 from 2015 to 2019 (103.2 versus 29.1 per 1,000 women) while slightly increase from 2019 to 2020 (29.1 versus 32.9 per 1,000 women). In 2021, the rate decreases to 29.4. The concurrent receipt of contraceptives with an opioid prescription is 13% in 2021 compared to 14% in 2020. As discussed, alternative forms of birth control may be used however cannot be captured in this analysis.