

## Tennessee Behavioral Risk Factor Surveillance Survey 2011 State and Regional Weighted Data

**During the past 12 months, have you had either a seasonal flu shot or a seasonal flu vaccine that was sprayed in your nose?**

State and Regions	Total Respondents	Flu Shot in Past 12 Mos? <sup>1</sup>	Weighted Percent <sup>2</sup>	Population With Flu Shot <sup>3</sup>
<b>TENNESSEE <sup>4</sup></b>	5,531	2,951	44.3	2,000,165
<b>Northeast</b>	431	244	47.3	125,772
<b>East</b>	339	193	47.0	250,561
<b>Southeast</b>	342	155	38.5	90,585
<b>Upper Cumberland</b>	384	219	51.6	129,043
<b>Mid-Cumberland</b>	365	198	43.1	348,130
<b>South Central</b>	355	186	36.4	97,593
<b>Northwest</b>	379	201	44.3	80,860
<b>Southwest</b>	405	201	42.8	87,101
<b>Shelby</b>	401	206	43.5	267,959
<b>Davidson</b>	378	206	43.0	202,234
<b>Knox</b>	404	251	54.8	173,373
<b>Hamilton</b>	372	179	39.2	99,189
<b>Madison</b>	385	202	45.6	32,482
<b>Sullivan</b>	432	244	46.2	54,093

<sup>1</sup> Number of respondents who answered YES.

<sup>2</sup> Percent of adult population who received a flu shot during the past 12 months based on the survey.

<sup>3</sup> Estimated adult population who received a flu shot during the past 12 months based on the survey. Note: The sum of regional estimates in this column will not add up to statewide total.

<sup>4</sup> Statewide estimates include survey by cell phone and different weighting methodology.